



**QUEEN'S
UNIVERSITY
BELFAST**

QED in ultra---high laser fields: current experimental results and perspectives

Sarri, G. (2017). QED in ultra---high laser fields: current experimental results and perspectives. Abstract from Quantum Electronics (PQE) conference, Snowbird, Utah, January 2017, .

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.



**QUEEN'S
UNIVERSITY
BELFAST**

QED in ultra---high laser fields: current experimental results and perspectives

Sarri, G. (2017). QED in ultra---high laser fields: current experimental results and perspectives. In G. R. Welch , & F. A. Narducci (Eds.), *Journal of modern Optcis: Colloquium on the physics of quantum electronics* Taylor & Francis.

Published in:
Journal of modern Optcis

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

QED in ultra-high laser fields: current experimental results and perspectives

G. Sarri

*School of Mathematics and Physics, Queen's University Belfast, BT7 1NN,
Belfast, United Kingdom*

Quantum Electrodynamics (QED) represents thus far one of the greatest theoretical achievements in modern physics, able to elegantly combine electromagnetism, quantum mechanics, and special relativity into a unified theory. Its predictions have been tested to a high degree of precision in relatively low field intensities but, thus far, little is experimentally known about the behaviour of quantum systems in ultra-high fields. QED effects are dominant if these fields are comparable to the critical field (also known as the Schwinger field): $E_S \sim 1.3 \times 10^{18}$ V/m. At these field intensities, exotic phenomena will occur, such as stochastic photon emission [1], electron-positron pair production even in a classical vacuum [2], and strong radiation reaction [3].

Even though electromagnetic fields of this nature are expected to be present around massive astrophysical objects (see, for instance, [4]), experimental studies have been extremely limited, only exploiting strong crystalline fields [5] or ultra-relativistic electron beams in accelerators [6].




Fig. 1 Electron energies and laser amplitudes achievable by different laser systems: ELI-NP (red rectangle), Astra-Gemini (yellow), and TARANIS-X (green) compared with different physical regimes. The red dashed line corresponds to the critical field in the rest frame of the electron. The grey-shaded area shows regions of pure non-linear QED. The blue circle and the red triangle relate to the SLAC experiment [6], and our Previous Work, respectively.

conditions where electrons feel, in their rest frame, an electric field equal to $0.2 E_S$ (red triangle in Fig.1). This was achieved exploiting a head-on collision of an intense laser pulse ($I_L \sim 2 \times 10^{20}$ Wcm⁻²) with a laser-driven ultra-relativistic electron beam (maximum Lorentz factor of $\gamma_L \sim 3500$). Clear evidence was found, for the first time, of strong radiation reaction with a 40% electron energy loss in 40 femtoseconds of laser duration [8]. This talk will present these results, together with a discussion of current developments in order to fully access experimentally, for the first time, regimes of purely non-linear QED.

References

- | | |
|---|---|
| [1] T. G. Blackburn et al. PRL 112, 015001 (2014) | [6] C. Bula et al., PRL 76, 3116 (1996) |
| [2] A. R. Bell et al., PRL 101, 200403 (2008). | [7] E. Esarey et al., Rev. Mod. Phys. 81, 1229 (2009) |
| [3] A. di Piazza et al, Rev. Mod. Phys. 84, 1177 (2012) | [8] G. Sarri et al. Nat. Phys. <i>to be submitted</i> . |
| [4] J. Sultana et al. Astrophys. J. 779, 16 (2013) | |
| [5] H. D. Hansen et al., PRD, 69, 032001 (2004) | |